

Soybean seed [Glycine max L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil

by yuliarahmawatiz991 1

Submission date: 06-Jul-2022 04:19PM (UTC+1000)

Submission ID: 1867207688

File name: Jamilah_2020_IOP_Conf._Ser._Earth_Environ._Sci._497_012039.pdf (688.67K)

Word count: 3362

Character count: 17043

PAPER • OPEN ACCESS

8
Soybean seed [*Glycine max* L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil

6
To cite this article: J Jamilah et al 2020 IOP Conf. Ser.: Earth Environ. Sci. **497** 012039

View the [article online](#) for updates and enhancements.

You may also like

- 1
- [Where should we apply biochar?](#)
Hamze Dokoohaki, Fernando E Miguez, David Laird et al.
- [Going beyond the green: senesced vegetation material predicts basal area and biomass in remote sensing of tree cover conditions in an African tropical dry forest \(miombo woodland\) landscape](#)
Marc Mayes, John Mustard, Jerry Melillo et al.
- [The effect of citric acid and sodium bicarbonate concentration on the quality of effervescent of red ginger extract](#)
Giyatmi and D K Lingga



The Electrochemical Society
Advancing solid state & electrochemical science & technology

241st ECS Meeting

May 29 – June 2, 2022 Vancouver • BC • Canada
Abstract submission deadline: Dec 3, 2021

Connect. Engage. Champion. Empower. Accelerate.
We move science forward



Submit your abstract



Soybean seed [*Glycine max* L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil

J Jamilah^{1*}, N Irawan¹, A S Thesiwati¹, and M Ernita¹

¹Agriculture Faculty, Tamansiswa University, Jl. Tamansiswa No. 9 Padang, West Sumatera, Indonesia 25138

E-mail: jamilahfatika@gmail.com

Abstract. Soybean yields in Indonesia tend to be low, on average $<1.5 \text{ Mg ha}^{-1}$ and many of the needs are met through soybean imports. The purpose of this study was to determine the effect of seed coating and sodium bicarbonate on the growth and yield of soybeans in alluvial soils. The experiment was carried out at Alluvial soil in Padang, with an altitude of 10 m above sea level. The experiments were arranged in a completely randomized design in the factorial form. The first-factor was the application of coating material to soybean seeds consisting of 2 levels; uncoated and coated. The second factor was the application of sodium bicarbonate in the leaves of the plant consisting of 4 concentrations; 0%; 0.125%; 0.250% and 0.375%, with 3 replications. Data were analyzed statistically using F-tests 5% significance level. If the treatment had a significant effect, it will continue using the LSD test 5% significance level. The parameters of observations were made; plant height when the generative phase, the number of primary branches, the number of nodules, the number of pith pods, the weight of 100 seeds, dry weight of seeds per plot. The result showed that the application of coating materials on soybean seeds and 0.125% sodium bicarbonate sprayed on the leaves increase the growth and yield of soybean Anjasmoro variety with the highest seed yield was 1.90 Mg ha^{-1} .

1. Introduction

Soybean production in Indonesia is still low at an average of 1.5 Mg ha^{-1} [1]. The original habitat of soybean crops. Soybeans will grow well in temperate or sub-tropical climates [2] stating that low soybean yields are also caused by one of them because technological innovation is still low. Nowadays, Indonesia imports soybeans 70% of the soybean demand, and the rest is produced domestically.

One of the simple technologies developed is making a coat for soybean seeds using ex soybean's soil, manure enriched with nutrients P and K. Results of the study [3] had proven that giving coated [1kg of topsoil + 1kg compost + 100 g lime + 0.5 kg of ex soybean's soil + 10 g PK fertilizer] increase soybean yield per plant not accompanied by the application of Unitas Super liquid organic fertilizer, and vice versa soybean crops that get a positive response to POC will grow well if the seeds are not coated. In this experiment, coated material was made no longer using topsoil and lime, but only a mixture of former soybean soil, manure, SP36, and KCl fertilizer. The possibility of the results could be better than previous experiments and the response of liquid fertilizer was also accompanied by coated seed. It is expected that with the application of the coating, the seeds come into direct contact with bacteria originating from soybean former soils, get complete nutrition, sufficient oxygen, thereby increasing the ability to grow seeds in alluvial soils.

² The application of organic liquid fertilizer derived from organic waste materials has been successfully applied to various plants, including; pruned lowland rice [4-6]. In general, they proved that the organic liquid fertilizer has a very positive effect on increasing crop yield, although the recommended concentration of application is not the same. Another effort to increase the fertility of soybean plants is by spraying sodium bicarbonate [baking soda] into the leaves of the plant. According to [7] giving 1 tablespoon of baking soda mixed with 2.5 tablespoons of oil dissolved in 1 gallon of water sprayed on the leaves of plants is good for horticultural plant growth. Sodium bicarbonate is effectively used as a control of fungal diseases in plants, black spots, powdery mildew, but must be careful in large amounts of use can have a negative effect on growth and yield. [8] Explained that by increasing the concentration of application from 0 to 0.35% sodium bicarbonate, sodium, sodium carbonate, sodium chloride and sodium sulfate to soil influence to reduce the growth of wheat germination. If the limit of tolerance for the wheat seedling is taken as the point of concentration when both germination and growth are prevented, this is found to be with the carbonate 0.13 percent, with sodium chloride 0.52 percent, and with sodium sulfate 0.56 percent. It is not clear why there was so little difference in these experiments between the limit of tolerance for sodium chloride and sodium sulfate. However, it is still unknown if sodium bicarbonate that was sprayed on the leaves can act as a substitute for nutrient needs, protect plants from pest and disease disorders or have a negative effect on soybean plants. The purpose of this study was to determine the effect of seed coating and sodium bicarbonate application on the growth and yield of soybeans in alluvial soils.

2. Materials and Methods

2.1. Location and plant materials

Experiments have been carried out in Alluvial, Kalumbuk village, Padang City with altitude at 10 m above sea level. Some materials used in this experiment included; The soybean seeds used are Anjasmoro varieties, coating materials, and sodium bicarbonate. The seed coating material is formed from the mixture 49% ex soybean's soil: 49% composted manure: 1% Potassium: 1% Phosphorus. All materials in a dry air condition are then mashed and sieved at a 2 mm diameter sift, evenly mixed which is called a coating material. Application of coating materials and seed by a ratio 1: 1.

Article Error (ETS)

Frag. (ETS)

2.2. Experimental design

The experiments were arranged in a completely randomized design in the factorial form. The first-factor was the application of coating material to soybean seeds consisting of 2 levels; without being coated and coated. The second factor was the application of sodium bicarbonate in the leaves of the plant consisting of 4 concentrations; 0%; 0.125%; 0.250% and 0.375%, with 3 replications. Data were analyzed statistically using F tests 5% significance level. If the treatment had a significant effect, so it will continue using the LSD test 5% significance level. Soybeans planted with a spacing of 30 x 15 cm with a plot size of 150 cm x 150 cm. Basic fertilizer given for each hectare was 25 kg of Urea; 150 kg SP36 and 100 kg KCl.

S/V (ETS)

2.3. Procedures and observations

Application of coating materials to soybean seeds by soaking the soybean seeds for 1 night, then the seeds coated with coating materials according to the treatment described above. The coating was applied until all ingredients evenly attached to the seed. Planting is done as much as 1 seed per planting hole as deep as 3 cm. The application of sodium bicarbonate starts at 2nd weeks after planting the seeds every 2 weeks and stops until the plants achieve the generative phase. The parameters of observations made included; plant height when the generative phase, the number of primary branches, the number of nodules, the number of pith pods, the weight of 100 seeds, dry weight of seeds per plot.

Article Error (ETS)

3. Results and Discussion

3.1. The effect of soybean seed coating and sodium on plant growth

The results of the experiment showed that there was a single effect of each of the soybean seed coating material and the concentration of sodium bicarbonate on height, but did not significantly affect the number of primary branches of the plant at 45 days after planting. The interaction of coating and sodium bicarbonate is seen in the formation of nodules, presented in Table 1.

Table 1. The effect of coating on soybean seeds and sodium bicarbonate on plant growth

Plant height at 45 days after planting [cm]					
treatments	Sodium bicarbonate concentration [%]				
Seed treatments	0	0.125	0.250	0.375	Average
uncoated	88.22	94.78	95.00	96.44	93.61 b
coated	97.67	98.44	99.44	99.44	98.75 a
Average	92.95 b	96.61 a	97.22 a	97.94 a	
CV [%]	2,68				
the number of primary branches per plant					
Treatments	Sodium bicarbonate concentration [%]				
Seed treatments	0	0.12	0.25	0.37	Average
uncoated	4.33	3.70	4.00	3.33	3.83
coated	3.33	3.67	4.00	4.00	3.75
Average	3.83	3.67	4.00	3.67	
CV[%]	15.23				
the number of nodules per plant					
Treatments	Sodium bicarbonate concentration [%]				
Seed treatments	0	0.125	0.250	0.375	Average
uncoated	6.67 Bb	7.00 Bb	9.00 Aa	8.33 Aab	
coated	10.67 Aa	10.00 Aa	8.00 Ab	9.00 Aab	
Average					
CV [%]	15.41				

The numbers followed by the same uppercase letters are not significant in the column and the numbers followed by the same lowercase letters are not significantly different in the lines according to the LSD level of 5%.

Plant growth was better in the treatment given coated than those not given coated. This shows that coated has a positive role in the seed from the germination process to high growth to the generative phase. Coating given is a nutritious material so that the seeds get enough food starting from the early budding phase to the mature phase. Adequate nutrition which obtained from manure and elements of P and K which are formulated in the form of coating preparations has provided sufficient nutrition for

soybean plants. It was also explained by [9] that at the laboratory scale seeds that were treated with lime material or other nutritional promote better germination growth than seeds uncoated.

In general, the application of sodium bicarbonate sprayed into the plant had a positive effect on plant height, but each concentration given from 0.125% to 0.375% was not significantly different. Sodium bicarbonate contains sodium and bicarbonate. Sodium plays a role in replacing the position of potassium in metabolism because of the same prevalent one. The general role of sodium in low concentrations can be an enzyme activator as well as, the same role as the K element [10]. They explained that potassium and sodium are both involved in enzyme activity. [11] explains that fertilization through leaves is more effective than giving it through the soil. The penetration of nutrients is through the surface leaf layers and their uptake across the plasma membrane of the epidermal cells. Also, environmental factors, aspects of plant biology and solution properties had a crucial effect on the efficiency of foliar fertilization, are presented. In general, the movement of low molecular weight solutes [organic acids, amino acids, sugars] from the leaf surface to the cell wall of the epidermis is a non-metabolic process that is driven by diffusion and electrochemical potential formed by an increase in negative charge across the cuticle membrane.

There was no significant effect of the seed coating treatment or the application of sodium bicarbonate to soybean leaves on the number of primary branches. The nature of the formation of primary branches is more dominated by the genetics of these plants. In general, Anjasmoro varieties produce several primary branches ranging from 2.9 and 5.6 [12].

The formation of nodules was influenced by salute and sodium bicarbonate preparations. Giving coated had a direct impact on the formation of nodules. Root nodules enhanced 50-80% at seed coated compared to an uncoated seed. The same thing had been proven by [3] that the formation of nodules was strongly influenced by the inoculation of *Rhizobium bacteria* in the form of legum or using soybean soil. Likewise, previously [13] reported that the presence of Rhizobacteria in the soil, enhance N fixation from the air to reduce the need for N nutrients and enhance soybean yields. The number of root nodules decreases as the concentration of sodium bicarbonate application increases at seed coated, but this does not occur in uncoated seeds. Negative effects begin to appear if the sodium bicarbonate concentration was increased to 0.250%. The negative effects of sodium application had also been explained [14] on wheat crops, and [8] sugar beet plants. However, these impacts vary depending on the type of plant, soil, and environment.

3.2. The effect of coating on soybean seeds and sodium bicarbonate on soybean yield components

Coating and sodium bicarbonate application did not have a significant effect on the number of seeds per pod, weighing 100 seeds, but there were interactions between the two treatments on seed yield per hectare, presented in Table 2.

Giving coated materials on seeds and 0.125% sodium bicarbonate gave the highest yield of soybean to the dry weight at 14% water content, but it was not significantly different at the same concentration uncoated. The results of this experiment proved that coating is important so that soybean plants become more vigorous and healthier so that they can produce high. Likewise, the application of sodium bicarbonate every 2 weeks until it reached the generative phase had a positive effect on the formation of dried soybeans. [2] which stated that soybean yield did not reach 2 Mg ha^{-1} , still could not be improved in this experiment. Whenever, compared to plants that were not coated, the application of a coating to soybean seeds only enhanced 1%. This due to the fact that the alluvial soils tested are indeed very low in organic matter content which characterized as bright yellow and the soil tends to be compact. The condition of lack of organic matter in the soil also affects the growth and development of *Rhizobium bacteria* in the soil so that it has an impact on yields that increase significantly [15]; explained that the diversity of microorganisms is largely determined by the chemical and physical properties of the soil.

Acidic and compact soil will reduce the presence of N-fixing bacteria so that the element is not easily accessed by the roots. The results of this experiment were still higher than the results [13] which obtained the dry seeds of Willis variety only $1,026 \text{ Mg ha}^{-1}$ from 10 Mg ha^{-1} compost and 100 kg ha^{-1}

nitrogen fertilizer application. According to the descriptions of Wilis and Anjasmoro varieties, the yield potential is almost the same.

Table 2. The effect of coating on soybean seeds and sodium bicarbonate on soybean yield components

The number of pith pods per plant					
Seed treatments	Sodium bicarbonate concentration [%]				
	0	0.125	0.250	0.375	Average
uncoated	109.44	108.22	113.44	109.89	110.25
coated	101.11	114.22	98.00	98.56	102.97
Average	105.28	111.22	105.72	104.22	
CC [%]	10.75				
the weight of 100 seeds					
Seed treatments	Sodium bicarbonate concentration [%]				
	0	0.125	0.250	0.375	Average
uncoated	9.63	9.42	10.45	8.74	9.56
coated	9.25	10.55	9.21	9.49	9.63
Average	9.44	9.98	9.83	9.11	
CC [%]	9.73				
the dry weight of seeds per hectare [Mg]					
Seed treatments	Sodium bicarbonate concentration [%]				
	0	0.125	0.250	0.375	Average
uncoated	1.75Aab	1.88Aa	1.85Aa	1.57 Bab	1.76
coated	1.78 Aab	1.90Aa	1.35Bc	1.66 Abc	1.67
Average	1.76	1.89	1.60	1.61	
CC [%]	9.66				

The numbers followed by the same uppercase letters are not significant in the column and the numbers followed by the same lowercase letters are not significantly different in the lines according to the LSD level of 5%.

4. Conclusion

Application of coating materials on soybean seed and 0.125% sodium bicarbonate sprayed on the leaves increase the growth and yield of soybean Anjasmoro variety with the highest dry seed was 1.90 Mg ha⁻¹.

References

- [1] BPS 2018 *Hasil kedelai Indonesia dalam angka* <https://www.bps.go.id/subject/53/tanaman-pangan.html#subjekViewTab3>.
- [2] Satria 2015 *Produksi Kedelai Nasional Masih Rendah* Berita UGM <https://ugm.ac.id/id/berita/9987-produksi-kedelai-nasional-masih-rendah>.
- [3] Erlinda, Jamilah, and Herman, W 2019 Pengaruh Sediaan Salut dan Pupuk Organik Cair Terhadap Pertumbuhan dan Hasil Kedelai (*Glycine max* L) *Solum*, **XVI** (1) 40–48

- [4] Jamilah 2017 Pengaruh Pupuk Organik Cair Asal *C.odorata* Terhadap Serapan Hara Kalium dan Hasil Padi Ladang *Jurnal Bibiet* **1** (1) 17–26
- [5] Jamilah and Mulyani SJ 2017 The Application of Liquid Organic Fertilizer of *Chromolaena odorata* on Ratooned Rice Plants Cultivation *As. J. of Appl. Res. for Com. Dvlp. and Emp.* **1** (1) 9–15
- [6] Jamilah, Soleh R, and Herman W 2017 Respon Tanaman Padi (*Oryza sativa* L.) Kabir 07 Terhadap Pupuk Organik Cair Crocober Plus Khusus Kota Padang dengan Iklim *Af. Jurnal Solum* **7** (1) 18–27
- [7] Baker and Marissa 2010 Effects of Sodium Bicarbonate on Plant Growth. Hunker <https://www.hunker.com/12236139/effects-of-sodium-bicarbonate-baking-soda-on-plant-growth>
- [8] Gazette SB and Mar N 1955 Influence of Sodium Bicarbonate on the Growth and Chlorosis of Garden Beets **116** (3) 201–209
- [9] James MS 2015 Effects of Biochar-Based Seed Coatings on Seed Germination and Seedling Vigor of California Brome (*Bromus carinatus* L.) and Blue Wildrye (*Elymus glaucus* L.) Abstract Oregon
- [10] Mengel K and Kirkby 2001 *Principles of plant nutrition* 5th eds Kluwer Academic Publishers <https://doi.org/10.1093/aob/mch063>
- [11] Wójcik P 2004 Uptake of mineral nutrients from foliar fertilization. *Journal of Fruit and Ornamental Plant Research* **12** (Special ed.: Orchard management in sustainable fruit production) 201–218
- [12] Irawan N 2019 Pengaruh sediaan salut an pemberian baking soda terhadap pertumbuhan dan hasil kedelai Undergraduate Thesis-Fakultas Pertanian Universitas Tamansiswa Padang
- [13] Tridiati, Mubarik NR and Rasmata Y 2013 Respon Pertumbuhan Tanaman Kedelai terhadap *Bradyrhizobium japonicum* Toleran Masam dan Pemberian Pupuk di Tanah Masam. *J. Agron. Indonesia* **41** (1) 24–31
- [14] Barrow NJ and TC Shaw 1978 Sodium bicarbonate as an extractant for soil phosphate III. Effects of the buffering capacity of soil for phosphate. *Geoderma* **16** (4) 273-283
- [15] Fernandes EG, OL Pereira, CC Silva, CBP da Bento, and MV de Queiroz 2015 Diversity of endophytic fungi in *Glycine max*. *Micro. Res.* **181** 84-92

Acknowledgment

Thank you to the dean of Agriculture Faculty of Tamansiswa University who facilitates all these activities and the students who helped.

Soybean seed [Glycine max L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil

ORIGINALITY REPORT

98%
SIMILARITY INDEX

42%
INTERNET SOURCES

98%
PUBLICATIONS

92%
STUDENT PAPERS

PRIMARY SOURCES

1 J Jamilah, N Irawan, A S Thesiwati, M Ernita. "Soybean seed [Glycine max L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil", IOP Conference Series: Earth and Environmental Science, 2020
Publication **67%**

2 Submitted to Universitas Andalas
Student Paper **23%**

3 Submitted to Universitas Bangka Belitung
Student Paper **5%**

4 Nasruddin, A T Bondan, S Agustini. "The effect of sodium bicarbonate ratio for the mechanical properties of underarm pads rubber for crutches", IOP Conference Series: Materials Science and Engineering, 2020
Publication **1%**

5 darchive.mblwhoilibrary.org
Internet Source **1%**

6 www.coursehero.com
Internet Source **<1%**

7

Submitted to Aberystwyth University

Student Paper

<1 %

8

iopscience.iop.org

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Soybean seed [Glycine max L.] coated by fertile soil-applied sodium bicarbonate at alluvial soil

PAGE 1

PAGE 2



Prep. You may be using the wrong preposition.



Confused You have used **A** in this sentence. You may need to use **an** instead.



Prep. You may be using the wrong preposition.



S/V This subject and verb may not agree. Proofread the sentence to make sure the subject agrees with the verb.



Article Error You may need to use an article before this word.



Frag. This sentence may be a fragment or may have incorrect punctuation. Proofread the sentence to be sure that it has correct punctuation and that it has an independent clause with a complete subject and predicate.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Proofread This part of the sentence contains a grammatical error or misspelled word that makes your meaning unclear.



Wrong Article You may have used the wrong article or pronoun. Proofread the sentence to make sure that the article or pronoun agrees with the word it describes.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Article Error You may need to remove this article.



Article Error You may need to use an article before this word.



Missing ",," You may need to place a comma after this word.

PAGE 3



Article Error You may need to use an article before this word. Consider using the article **the**.



Article Error You may need to use an article before this word.



Article Error You may need to use an article before this word. Consider using the article **the**.



Article Error You may need to use an article before this word.



Article Error You may need to remove this article.



Article Error You may need to remove this article.



Proofread This part of the sentence contains a grammatical error or misspelled word that makes your meaning unclear.



Article Error You may need to use an article before this word.



Frag. This sentence may be a fragment or may have incorrect punctuation. Proofread the sentence to be sure that it has correct punctuation and that it has an independent clause with a complete subject and predicate.



Article Error You may need to use an article before this word.



S/V This subject and verb may not agree. Proofread the sentence to make sure the subject agrees with the verb.



Confused You have used **hole** in this sentence. You may need to use **whole** instead.



Article Error You may need to use an article before this word.



Missing "," You may need to place a comma after this word.



Missing "," You may need to place a comma after this word.



Confused You have used **a** in this sentence. You may need to use **an** instead.



Missing "," You may need to place a comma after this word.



Article Error You may need to use an article before this word.



Article Error You may need to use an article before this word. Consider using the article **the**.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Article Error You may need to use an article before this word. Consider using the article **the**.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Article Error You may need to remove this article.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Article Error You may need to use an article before this word.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Missing "," You may need to place a comma after this word.



Article Error You may need to use an article before this word.



Missing "," You may need to place a comma after this word.



S/V This subject and verb may not agree. Proofread the sentence to make sure the subject agrees with the verb.